

OLIGOMERIC FLAVANOIDS. PART 15^a. BASE-CATALYZED PYRAN REARRANGEMENTS OF
PROCYANIDIN B-2, AND EVIDENCE FOR THE OXIDATIVE TRANSFORMATION OF B- TO
A-TYPE PROCYANIDINS

JOHANN F.W. BURGER^b, HERBERT KOLODZIEJ^b, RICHARD W. HEMINGWAY^c, JAN P.
STEYNBERG^b, DESMOND A. YOUNG^b, AND DANEEL FERREIRA^{b*}.

^b Department of Chemistry, University of the Orange Free State, P.O. Box 339
Bloemfontein, 9300 South Africa

^c Southern Forest Experiment Station, USDA-Forest Service, 2500
Shreveport Highway, Pineville, Louisiana, 71360 U.S.A.


(Received in UK 30 April 1990)

Abstract — Procyanidin B-2 **3** is subject to facile C-ring isomerizations in 0.1M NaHCO₃ solution to form a novel series of 3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-*h*]chromenes **7**, **9**, and **10**. The low percentage conversion of B- to A-type procyanidin **2** is rationalized in terms of an initial oxidative removal of hydride ion at C-2 (C-ring).

Proanthocyanidin A-2 **1** [(*-*)-epicatechin(4 β \rightarrow 8, 2 β \rightarrow O \rightarrow 7)-(*-*)epicatechin] was first isolated by Mayer *et al.* from the seed of *Aesculus hippocastanum*¹. The structure was deduced by Haslam and his co-workers *via* spectroscopic and chemical evidence² and has, more recently, been unequivocally established by X-ray crystallography³. A variety of proanthocyanidins possessing the doubly-linked unit of type **1** has since been reported^{2,4-10}. Owing to the close structural relationship between proanthocyanidin A-2 **1** and procyanidin B-2 **3**, Porter¹¹ has proposed a biosynthetic pathway for the conversion of B- to A-type procyanidins which involves an enzyme mediated hydroxylation at C-2 (C-ring) of **3**. Despite the considerable progress in the semi-synthetic approach towards condensed tannins over the last decade similar efforts for the A-type procyanidins are limited to a fortuitous reference⁹ to the oxidative conversion of procyanidin B-1 to proanthocyanidin A-1, but with no experimental details. Our recent investigations of the base-catalyzed pyran rearrangements of proflisetinidins^{12,13} and procyanidin B-3¹⁴ which emphasizes the involvement of 7-OH(D) and C-2(C) (*cf.* structure **3**), in conjunction with the industrial interest^{15,16} in the base-simulated reactions of procyanidins, prompted application of a similar protocol to procyanidin B-2 **3**.

Owing to the lability of the interflavanyl bond in procyanidins at alkaline pH^{14,17}, the conditions previously¹²⁻¹⁴ employed were adjusted for procyanidin B-2. Thus, treatment of (4 β ,8)-bis-(*-*)

*Part 14. J.C.S. Malan, P.J. Steynberg, J.P. Steynberg, D.A. Young, B.C.B. Bezuidenhout, and D. Ferreira, Tetrahedron, 1990, paper AM 9409(2)



Scheme: Proposed routes to the formation of phlobatannins 7, 9, 10, and the C-2(F) epimer 2 (A-4) of procyanidin A-2

epicatechin **3** with 0.1M NaHCO₃ (pH 8.15) for 6.5 h at 40°C under nitrogen containing traces of oxygen, gave complete conversion into a mixture consisting of oligomeric procyanidins (ca. 70%) and five mobile fractions following chromatography on Sephadex LH-20 in ethanol. The latter fractions afforded a mixture (47:53 by ¹H NMR analysis) of (-)-epicatechin **4** and (-)-catechin **5** (J_{2,3} ca. 1.0 and 8.0 Hz respectively), and four pure compounds with modified C-rings (Scheme). These comprised the functionalized 8,9-*cis*-9,10-*trans*-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-*h*]chromene **9** as the major product, its C-2(F) epimer **10**, the 8,9-*trans*-9,10-*cis* analogue **7**, and the C-2(F) epimer **2** of proanthocyanidin A-2 **1**. The relative ease of purification of these phenols contrasts with those derived from the proflisetinidins^{12,13} where the additional chromatographic stages offered by successive methylation and acetylation were a prerequisite for compound purity.

The ¹H NMR spectra of the tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10** are conspicuously free of the effects of dynamic rotational isomerism¹⁸ at ambient temperatures hence indicating their pyran rearranged nature¹². ¹H NMR coupling constants (Table) of the heterocyclic proton resonances (J_{2,3} ca. 1.0 Hz for **7** and **9**, J_{2,3} 8.5 Hz for **10**; J_{8,9} 7.5, J_{9,10} 5.0 Hz for **7**; J_{8,9} ca. 1.0 Hz for **9** and **10**, J_{9,10} 2.5 and 2.0 Hz for **9** and **10** respectively) are in accordance¹² with 2,3-*cis*-8,9-*trans*-9,10-*cis* relative configuration for **7**, 2,3-*cis*-8,9-*cis*-9,10-*trans* for **9**, and 2,3-*trans*-8,9-*cis*-9,10-*trans* for **10**.

Table ¹H NMR peaks of the tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10**, and the A-type procyanidin **2** in (CD₃)₂CO (23°C) at 300 MHz. Splitting patterns and J-values (Hz) are given in parentheses.

Ring	Proton	7	9	10	2
A	3/5 or 6/8	5.98(br.s)	5.90(d,2.0) 5.98(d,2.0)	5.94(d,2.0) 6.01(d,2.0)	6.08(d,2.5) 6.10(d,2.5)
B	2	6.86(d,2.0)	6.65(d,2.0)	6.63(d,2.0)	7.17(d,2.0)
	5	6.74(d,8.0)	6.70(d,8.0)	6.68(d,8.0)	6.81(d,8.0)
	6	6.67(dd,2.0,8.0)	6.46(dd,2.0,8.0)	6.47(dd,2.0,8.0)	7.03(dd,2.0,8.0)
	8	4.99(d,7.5)	5.65(br.s,ca.1.0)	5.59(br.s,ca.1.0)	3-H: 4.16(d,3.5)
	9	4.21(dd,5.0,7.5)	4.30(dd,1.0,2.5)	4.15(dd,1.0,2.0)	4-H: 4.29(d,3.5)
	10	4.73(d,5.0)	4.22(d,2.5)	4.19(d,2.0)	
D	6	6.06(s)	6.00(s)	5.86(s)	5.97(s)
E	2	6.57(d,2.0)	6.66(d,2.0)	6.85(d,2.0)	6.83(d,2.0)
	5	6.58(d,8.0)	6.77(d,8.0)	6.76(d,8.0)	6.73(d,8.0)
	6	6.17(dd,2.0,8.0)	6.12(dd,2.0,8.0)	6.69(dd,2.0,8.0)	6.67(dd,2.0,8.0)
F	2	4.76(br.s,ca.1.0)	4.69(br.s,ca.1.0)	4.53(d,8.5)	4.56(d,7.5)
	3	4.07(m)	4.12(m)	3.97(m)	3.95(m)
	4	2.70(dd,3.0,17.0)	2.69(dd,2.5,17.0)	2.49(dd,8.0,16.0)	2.54(dd,8.0,17.0)
	4 _{ax}				
	4 _{eq}	2.81(dd,4.5,17.0)	2.85(dd,5.0,17.0)	2.83(dd,5.5,16.0)	2.89(dd,5.5,17.0)

In contrast to the analogous pyran rearranged products from the base-catalyzed conversion of procyanidin

B-3¹⁴ which were characterized as octamethyl ether diacetates thus facilitating demonstration of a 'liberated' phloroglucinol A-ring by ¹H nuclear Overhauser effect (n.O.e.) difference spectroscopy, a different approach had to be adopted towards the structural elucidation of free phenols **7**, **9**, and **10**. The ¹H NMR spin systems in all three analogues were differentiated and fully analyzed by extensive spin-decoupling experiments using the benzylic 2-, 8-, and 10-H resonances as reference signals. A weak but structurally significant n.O.e. association (0.9, 1.9% resp.)^b of the broadened two-proton singlet (δ5.98) of the A-ring with 2- and 6-H(E) (δ6.57, 6.17 resp.) not only confirmed the tetrahydropyrano[2,3-*h*]chromene arrangement for analogue **7** but also established the location of the phloroglucinol moiety at C-10. It furthermore indicated a *cis*-relationship¹³ of the aryl substituents at C-2 and -10. When taken in conjunction with the known α -orientation of the E-ring of the (-)-epicatechin DEF unit, this also indicated an α -orientation for the phloroglucinol A-ring. A strong negative Cotton effect (CE) at 237 nm in the CD spectrum confirmed the 10 α -aryl substituent by application of the aromatic quadrant rule¹⁹. These CD and ¹H NMR (Table) features collectively facilitated definition of the absolute configuration of **7** as 2*R*,3*R*:8*S*,9*R*,10*R*.

The spin-decoupling experiments on analogues **9** and **10** also established a benzylic connection of 2-H(F) (δ4.69, 4.53 for **9** and **10** resp.) and 10-H(C) (δ4.22, 4.19 for **9** and **10** resp.) with the 2- and 6-protons of the respective pyrocatechol E- and B-rings. Besides the n.O.e. effect of 8-H(C) (δ5.65, 5.59 for **9** and **10** resp.) with 2- and 6-H (3.5, 3.8 and 3.9, 4.3% for **9** and **10** resp.) of a pyrocatechol moiety in both **9** and **10**, reminiscent of these structural types with a 3,4-disubstituted aryl group at C-10¹², the former proton did not exhibit additional long-range coupling thus reflecting the presence of an *ortho*-disubstituted phenyl residue at C-10. These observations hence implied an 'interchange' of the phloroglucinol A- and pyrocatechol B-rings in analogues **9** and **10** relative to the positions of these rings in the tetrahydropyrano[2,3-*h*]chromene **7c**. The chemical shifts of 8- and 10-H(C) and thus unambiguous proof for such an A-/B-ring interchange in **9** were confirmed by 2D-heteronuclear correlation of these protons with, respectively, C-8 and 10- (δ69.04, 45.33 resp.). This experiment also facilitated assignment of the chemical shifts of all H-bearing carbons (*cf.* Experimental). Confirmation of both the *cis*-relationship between the C-2 and -10 pyrocatechol E- and B-rings, both α -orientated (*vide supra*), and the tetrahydropyrano[2,3-*h*]chromene arrangement of **9** was again furnished by the n.O.e. associations of both 5- and 6-H(B) (δ6.70, 6.46 resp.) with 6-H(E) (δ6.12) (0.7, 1.1% resp.). A high-amplitude negative CE at 238 nm in the CD spectrum confirmed the 10 α -aryl substituent hence facilitating, in conjunction with ¹H NMR coupling constants of C-ring protons, definition of 2*R*,3*R*:8*S*,9*S*,10*R* absolute configuration for **9**. Although insufficient sample quantities precluded similar confirmation of the chemical shifts of 8- and 10-H(C) in **10**, the very similar shifts of the C-ring protons in **9** and **10** (Table) were taken as sufficient proof of a [2,3-*h*]chromene arrangement for **10**. These small chemical shift differences may be attributed to the β -orientated E-ring in the latter compound. The conspicuous absence of n.O.e. associations between the protons of the pyro-

^bApproximated value owing to signal overlap

^cCompare refs. 12-14 for similar phenomena occurring during base-catalyzed conversions of proflisetinidins and procyanidin B-3

catechol B- and E-rings in **10** presumably indicated a *trans*-relationship of these rings and hence a 10α B-ring. The CD spectrum, however, exhibited an intense positive CE at 239 nm which apparently reflected a 10β B-ring. Such a contradiction was commonly encountered for profisetinidin related analogues^{12,13} with 2,10-*trans* aryl groups and had been explained by significant contributions of A-conformers^{20,21} (F-ring) reversing the sign of the low-wavelength CE. We thus favour the 2*S*,3*R*:8*S*,9*S*,10*R* absolute configuration depicted for analogue **10**. The inversion of the absolute configuration at C-9 in **9** and **10** relative to that at C-3 in the biflavanoid precursor **3** is explained later.

The ¹H NMR spectrum (Table) of the remaining analogue **2** displayed in the heterocyclic proton region an AB-system [δ 4.16 (3-H), 4.29 (4-H), $J_{3,4}$ 3.5 Hz] characteristic² of the C-ring protons of A-type procyanidins. Confirmation of the chemical shift of 3-H was obtained by its pronounced n.o.e. association with both 2- and 6-H (B; 3.9, 4.3% resp.). Epimerization²³ at 2-C(F) under the mild basic conditions and hence conversion of the (-)-epicatechin DEF unit in precursor **3** to a (-)-catechin moiety in **2** was evident from the coupling constant ($J_{2,3}$ 7.5 Hz) of 2- and 3-H(F). A high-amplitude positive CE at 234 nm in the CD spectrum confirmed the β -orientation at C-4 and, combined with ¹H NMR data and known absolute configuration of procyanidin B-2 **3**, subsequently also the configuration as is depicted in **2**. Analogue **2** therefore represents the C-2(F) epimer of procyanidin A-2 which was previously isolated by Nishioka *et al.*⁹ and designated proanthocyanidin A-4.

Under the mild basic conditions procyanidin B-2 **3** is presumably transformed to an intermediate B-ring quinone-methide²² **6**^d which then serves as common precursor to the novel tetrahydropyrano[2,3-*h*]-chromenes **7**, **9**, and **10**. Analogue **7** originates *via* stereospecific pyran recyclization¹²⁻¹⁴ involving 7-OH(D) and the *si*-face at C-2 in quinone-methide **6**. Migration of the (-)-epicatechin moiety, assisted by the strongly electron-releasing phloroglucinol unit at C-4, to the *si*-face at C-2 in **6** and subsequent recyclization *via* 7-OH(D) and the *si*-face at C-4 in quinone-methide **6**^d may feasibly rationalize the genesis of the tetrahydropyrano[2,3-*h*]chromene with its 'interchanged' phloroglucinol A- and pyrocatechol B-rings. Inversion of configuration²¹ at the equivalent of C-3(C) in procyanidin B-2 **3** associated with the observed ring interchange is substantiated by CD data (*vide supra*). The susceptibility of the E-ring in *eg.* **3**, **6**, and **8** to quinone-methide formation at alkaline pH^{12-14,22} presumably also initiates epimerization at C-2(F) and hence formation of the (-)-catechin DEF unit in the ring-interchanged tetrahydropyrano[2,3-*h*]chromene **10**. Generation of the (-)-epicatechin/(-)-catechin mixture **4** and **5** is attributable to a similar phenomenon following cleavage of the base-labile¹⁷ interflavanyl bond in procyanidin B-2. A putative A-ring quinone-methide^{14,17} resulting from such a bond rupture may also induce the formation of the condensed analogues of unknown constitution *via* uncontrolled condensation with procyanidin B-2.

Formation of the A-/B-ring interchanged analogues **9** and **10**, arising *via* the exclusive 1,3-migration of the (-)-epicatechin moiety in quinone-methide **6**, contrasts with results for procyanidin B-3¹⁴ where pre-

^dQuinone-methides **6** and **8** are postulated and have not been isolated

ferential migration of the phloroglucinol A-ring at 4-C in an analogous quinone-methide was observed. Such an exclusivity in the shift of the (-)-epicatechin unit with its reduced migratory aptitude compared to that of the phloroglucinol moiety at C-4 for quinone-methide **6**, is presumably attributable to steric factors. The *cis*-coplanarity of the β -orientated sp^3 -orbital adjoining C-4 and C-8(D) and the electron-deficient *p*-orbital at the *si*-face of C-2 should provide the additional driving force for the 1,3-migration of the (-)-epicatechin unit in **6**.

The transformation of procyanidin B-2 **3** into the C-2(F) epimer **2** of procyanidin A-2 **1** presumably involves the oxidative removal of hydride ion at C-2(C) as the initial step. The nature of the oxidizing species is, however, not clear. Although the trace amounts of oxygen may effect the transformation **3** \rightarrow **11**, it seems more reasonable to suggest that the prevailing conditions induce oxidation of the *o*-dihydroxy functionality of the pyrocatechol B- or E-rings to an *o*-quinone²³ which subsequently serves as oxidant for the conversion **3** \rightarrow **11**. Experiments aimed at verifying the latter proposal *via* selective protection of the 3'-OH groups of the pyrocatechol rings of **3** and related procyanidins are presently being investigated. These results will be discussed in an impending publication. The results presented here nevertheless provide unequivocal chemical evidence in favour of the β -orientated doubly-linked unit of procyanidin A-2 **1** and of its C-2(F) epimer **2**. The apparently exclusive formation of procyanidin A-4 **2** is explicable in terms of formation of an E-ring quinone-methide of either **3** or **11** and subsequent recyclization to the thermodynamically more stable 2,3-*trans* relative configuration.

Under these extremely mild conditions we could not find evidence of rearrangements of procyanidin B-2 to catechinic acid-type products^{24,25} reputed for either decreasing its reactivity towards aldehydes or enhancing acidity^{24,26,27}. Our results, when considered in conjunction with similar observations for procyanidin B-3¹⁴, hence indicate that with the proper selection of conditions, extraction of conifer barks^{15,16} at alkaline pH may be performed without the adverse effects which have hitherto hampered the successful application of such an approach towards the economically important procyanidins. The 'liberated' phloroglucinol-type A-rings in tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10** should, indeed, lead to increased reactivity towards aldehydes compared to that of procyanidin B-2 **3**.

EXPERIMENTAL

¹H NMR spectra were recorded on a Bruker AM-300 spectrometer, for (CD₃)₂CO solutions (D₂O exchange) with Me₄Si as internal standard. Mass spectral data were obtained with a Kratos MS80 instrument, and CD data in MeOH on a Jasco J-20 spectropolarimeter. TLC was performed on precoated Merck plastic sheets (DC-Plastikfolien Kieselgel 60 PF₂₅₄, 0.25 mm) and compounds were located by H₂SO₄-HCHO (40:1 v/v) spray reagent. CC was done on Sephadex LH-20 in EtOH at a flow rate of 0.5 cm³ min⁻¹. Evaporations were done under reduced pressure at ca. 60°C in a rotary evaporator.

Base-catalyzed Conversion of (4/8)-Bis(-)-epicatechin **3.** — Procyanidin B-2²⁸ **3** (500 mg) was dissolved in 0.1M NaHCO₃ (200 cm³) and the mixture was stirred for 6.5 h at 40°C under nitrogen containing traces of oxygen. The mixture was chilled with crushed ice, acidified (0.1M HCl) to pH 6, and extracted with EtOAc (8x200 cm³). Drying (Na₂SO₄) of the extract followed by evaporation of the solvent afforded a brown, amorphous solid (330 mg). This was subjected to CC (2.5x90 cm column; 15 cm³/tube; first 400 cm³ of eluant discarded) to give the following fractions: 1(tubes 56-66, 32 mg), 2(114-132, 17 mg), 3(146-150, 6 mg), 4(160-180, 29 mg), 5(211-240, 15 mg), and 6(245-280, 63 mg).

Fraction 1 consisted of a (-)epicatechin/(-)catechin mixture (47:53) and fraction 6 consisted of high-molecular-mass analogues of procyanidin B-2. Owing to its complexity this mixture was not further investigated.

Fraction 2 afforded the C-2(F) epimer 2 (A-4) of procyanidin A-2 as a white solid (Found: M^+ , 576.1260. $C_{30}H_{24}O_{12}$ requires M , 576.1268); 1H NMR data (Table); CD $[\theta]_{297}$ 0, $[\theta]_{278}$ 1.0×10^4 , $[\theta]_{255}$ 0, $[\theta]_{234}$ 6.9×10^4 , and $[\theta]_{200}$ 1.9×10^4 .

Fraction 3 gave (2R,3R:8S,9R,10R)-3,5,9-trihydroxy-2,8-bis-(3,4-dihydroxyphenyl)-10-(2,4,6-trihydroxyphenyl)-2,3-cis-8,9-trans-9,10-cis-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 7 as a white solid (Found: M^+ , 578.1418. $C_{30}H_{26}O_{12}$ requires M , 578.1424); 1H NMR data (Table); CD $[\theta]_{320}$ 0, $[\theta]_{276}$ -5.3×10^4 , $[\theta]_{252}$ -1.4×10^4 , $[\theta]_{237}$ -8.9×10^4 , and $[\theta]_{226}$ 0.

Fraction 4 afforded (2R,3R:8S,9S,10R)-3,5,9-trihydroxy-2,10-bis-(3,4-dihydroxyphenyl)-8-(2,4,6-trihydroxyphenyl)-2,3-cis-8,9-cis-9,10-trans-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 9 as a white solid (Found: M^+ , 578.1431. $C_{30}H_{26}O_{12}$ requires M , 578.1424); 1H NMR data (Table); δ_c [(CD₃)₂CO; 23°C, 75.4 MHz] 96.07, 94.92 (C-3 + -5, A-ring), 116.27 (C-2, B)^e, 115.95 (C-5, B), 120.24 (C-6, B), 69.04 (C-8, C), 73.99 (C-9, C), 45.33 (C-10, C), 97.55 (C-6, D), 113.14 (C-2, E)^e, 116.05 (C-5, E), 118.28 (C-6, E), 79.31 (C-2, F), 66.79 (C-3, F), and 29.03 (C-4, F); CD $[\theta]_{319}$ 0, $[\theta]_{277}$ 2.6×10^4 , $[\theta]_{246}$ 0, $[\theta]_{238}$ -7.6×10^4 , and $[\theta]_{224}$ 0.

Fraction 5 consisted of (2S,3R:8S,9S,10R)-3,5,9-trihydroxy-2,10-bis-(3,4-dihydroxyphenyl)-8-(2,4,6-trihydroxyphenyl)-2,3-trans-8,9-cis-9,10-trans-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 10 as a white solid (Found: M^+ , 578.1429. $C_{30}H_{26}O_{12}$ requires M , 578.1424); 1H NMR data (Table); CD $[\theta]_{302}$ 0, $[\theta]_{270}$ -1.2×10^4 , $[\theta]_{257}$ 0, $[\theta]_{239}$ 5.6×10^4 , and $[\theta]_{202}$ 0.

Acknowledgements

Support by the Foundation for Research Development, C.S.I.R., Pretoria, the Sentrale Navorsingsfonds of this University, and the Marketing Committee, Wattle Bark Industry of South Africa, Pietermaritzburg is acknowledged.

REFERENCES

1. Mayer, W.; Goll, L.; Arndt, E.V.; Mannschreck, A. *Tetrahedron Lett.*, 1966, 429.
2. Jacques, D.; Haslam, E.; Bedford, G.R.; Greatbanks, D. *J. Chem. Soc., Perkin Trans. 1*, 1974, 2263.
3. Otsuka, H.; Fujioka, S.; Komiya, T.; Mizuta, E.; Takemoto, M. *Yakugaku Zasshi*, 1982, 102, 162.
4. Weinges, K.; Kaltenhauser, W.; Marx, H.-D.; Nader, E.; Perner, J.; Seiler, D. *Annalen*, 1968, 711, 184.
5. Hikino, H.; Shimoyama, N.; Kasahara, Y.; Takahashi, M.; Konno, C. *Heterocycles*, 1982, 19, 1381.
6. Hikino, H.; Takahashi, M.; Konno, C. *Tetrahedron Lett.*, 1982, 23, 673.
7. Ohigashi, H.; Minami, S.; Fukui, H.; Koshimizu, K.; Mizutani, F.; Sugiara, A.; Tomana, T. *Agric. Biol. Chem.*, 1982, 46, 2555.
8. Nonaka, G.; Morimoto, S.; Nishioka, I. *J. Chem. Soc., Perkin Trans. 1*, 1983, 2139.
9. Nonaka, G.; Morimoto, S.; Kinjo, J.; Nohara, T.; Nishioka, I. *Chem. Pharm. Bull.*, 1987, 35, 149.
10. Morimoto, S.; Nonaka, G.; Nishioka, I. *Chem. Pharm. Bull.*, 1988, 36, 33.
11. Porter, L.J. in *The Flavanoids - Advances in Research since 1980*, ed. Harborne, J.B., Chapman and Hall, London, 1988, 59.
12. Steynberg, J.P.; Burger, J.F.W.; Young, D.A.; Brandt, E.V.; Steenkamp, J.A.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1988, 3323, 3331.
13. Burger, J.F.W.; Steynberg, J.P.; Young, D.A.; Brandt, E.V.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1989, 671.
14. Steynberg, J.P.; Bezuidenhout, B.C.B.; Burger, J.F.W.; Young, D.A.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1990, 203.
15. Laks, P.; Hemingway, R.W. *J. Chem. Soc., Perkin Trans. 1*, 1987, 465, and refs. cited therein.

16. Laks, P.E.; Hemingway, R.W.; Conner, A.H. *J. Chem. Soc., Perkin Trans. 1*, 1987, 1875, and refs. cited therein.
17. Hemingway, R.W.; Foo, L.Y. *J. Chem. Soc., Chem. Commun.*, 1983, 1035.
18. Du Preez, I.C.; Rowan, A.C.; Roux, D.G. *Chem. Commun.*, 1971, 315, and refs. cited therein.
19. Van der Westhuizen, J.H.; Ferreira, D.; Roux, D.G. *J. Chem. Soc., Perkin Trans. 1*, 1981, 1220.
20. Porter, L.J.; Wong, R.Y.; Benson, M.; Chan, B.G. *J. Chem. Res.*, 1986, (S) 86; (M) 830.
21. Steynberg, J.P.; Burger, J.F.W.; Young, D.A.; Brandt, E.V.; Ferreira, D. *Heterocycles*, 1989, **28**, 923.
22. Whalley, W.B.; Mehta, P.P. *J. Chem. Soc.*, 1963, 5327.
23. Musso, H. in *Oxidative Coupling of Phenols*, eds. Taylor, W.I. and Battersby, A.R., Marcel Dekker, Inc., New York, 1967, 75.
24. Sears, K.D.; Casebier, R.L.; Hergert, H.L.; Stout, G.H.; McCandlish, L.E. *J. Org. Chem.*, 1974, **39**, 3244.
25. Courbat, P.; Weith, A.; Albert, A.; Pelter, A. *Helv. Chim. Acta*, 1977, **60**, 1665.
26. Herrick, F.H. *J. Agric. Food Chem.*, 1980, **28**, 228.
27. Kiatgrajai, P.; Wellons, J.D.; Gollub, L.; White, J.D. *J. Org. Chem.*, 1982, **47**, 2910.
28. Kolodziej, H.; Ferreira, D.; Roux, D.G. *J. Chem. Soc., Perkin Trans. 1*, 1984, 343.